

arXiv:hep-th/9404024v1 6 Apr 1994

A Parity Invariant Regularization in 3-D Quantum Electrodynamics

TADAHIKO KIMURA*

*Department of Physics, Faculty of Science, Chiba University
1-33 Yayoi-cho, Inage-ku, Chiba 263, Japan*

CHIBA-EP-76
hep-th/9404024
April 1994

Abstract

We regularize the 3-D quantum electrodynamics by a parity invariant Pauli-Villars regularization method. We find that in the perturbation theory the Chern-Simons term is not induced in the massless fermion case and induced in the massive fermion case.

*E-mail address: kimura@cuphd.nd.chiba-u.ac.jp

There has been much recent interest in Chern-Simons theories [1] in three dimensions in several contexts; their ability [2] to explain certain behaviour in fractional quantum Hall effect and high T_c superconductivity, their connection [3] with knots and conformal field theories in 2-D, and so on.

One of the important problems when studying Chern-Simons theories is the method employed to regularize ultraviolet divergent integrals. Three attempts have been made to regularize the theories; i) the Pauli-Villars regularization [4] [5], ii) the analytic regularization [6] and iii) the dimensional regularization [7] [8]. Unfortunately the results have not been totally conclusive, and all or some of these regularization methods seem to suffer from their own difficulties. One important such discrepancy which is concerned in this note is whether the Chern-Simons terms are induced through quantum fermion loop corrections even if the bare Lagrangian does not contain such terms. Until now it has been argued [4] [6] [8] that for massless fermions Chern-Simons term is induced in the Pauli-Villars regularization and not induced in the analytic and dimensional ones, and for massive fermions the Chern-Simons term is not induced in the Pauli-Villars regularization and induced in the analytic and dimensional ones. The reason why the Chern-Simons term is (is not) induced for massless (massive) fermions in the Pauli-Villars regularization method is the following. The usual Pauli-Villars regularization in 3-D breaks parity due to the regulator fermion mass and the massive regulator fermion loop induces the Chern-Simons term which remains finite even after the regulator masses tend to infinity. For massive fermion case this induced Chern-Simons term cancels the one induced through original fermion loop, and for massless fermion case it survives. Therefore it is desirable to develop a parity invariant Pauli-Villars regularization method in which the regulator fields do not induce any Chern-Simons term to discuss whether the Chern-Simons terms are induced through the quantum fermion loop correction or not.

On the other hand recently Frolov and Slavnov [9] has proposed a method of chiral gauge invariant regularization for the fermion field in the 16-component irreducible spinor representation of $SO(10)$ chiral gauge theory, which is automatically anomaly-free. The method is based on the introduction of infinite number of the Pauli-Villars regulator spinor fields which is in the 32-component reducible representation of $SO(10)$. The method seems to work for any anomaly-free chiral gauge theories [10].

The purpose of this short note is to clarify this problem in the perturbation theory by adapting a parity invariant Pauli-Villars regularization method which is a kind of 3-D variant of the chiral gauge invariant Pauli-Villars regularization proposed by Frolov and Slavnov [9]. Our conclusion is that this parity invariant Pauli-Villars regularization gives the same result as that in analytical and dimensional regularization methods.

We consider the 3-D quantum electrodynamics with one flavour of two-component spinor ψ of mass m . Note that one two-component spinor is counted as one flavour. In order to regularize the ultraviolet divergent integrals in parity invariant way, following Frolov and Slavnov, we must introduce the infinite number of fermionic and bosonic Pauli-Villars regulator spinor fields which belong to the 4-component reducible representation of the 3-D Lorentz group; ψ_r and ϕ_s , respectively. The fermions with the odd number of flavours cannot be regularized by the regulator fields with the finite even number of flavours. In the case of even number of flavours it is sufficient to introduce the finite number (a half number of flavours) of 4-component regulator spinor fields. With the two-component irreducible spinors they are written as $\psi_r = (\psi_{r+}, \psi_{r-})$ and $\phi_s = (\phi_{s+}, \phi_{s-})$. The parity transformation of the two-component spinors ψ , $\psi_{r\pm}$ and $\phi_{s\pm}$ is defined as

$$\begin{aligned} x = (x_0, x_1, x_2) &\rightarrow \bar{x} = (x_0, -x_1, x_2), \\ \psi(x) &\rightarrow \sigma_1 \psi(\bar{x}), \psi_{r\pm}(x) \rightarrow \sigma_1 \psi_{r\mp}(\bar{x}), \phi_{s\pm}(x) \rightarrow \sigma_1 \phi_{s\mp}(\bar{x}). \end{aligned} \quad (1)$$

Then the Lagrangian density regularized in a gauge and parity invariant way is given by

$$\begin{aligned} L = & -\frac{1}{4}F_{\mu\nu}^2 + \bar{\psi}i\gamma^\mu D_\mu \psi + m\bar{\psi}\psi \\ & + \sum_{r=1}^{\infty} \{ \bar{\psi}_{r+} i\gamma^\mu D_\mu \psi_{r+} + \bar{\psi}_{r-} i\gamma^\mu D_\mu \psi_{r-} + m_r (\bar{\psi}_{r+} \psi_{r+} - \bar{\psi}_{r-} \psi_{r-}) \} \\ & + \sum_{s=1}^{\infty} \{ \bar{\phi}_{s+} i\gamma^\mu D_\mu \phi_{s+} + \bar{\phi}_{s-} i\gamma^\mu D_\mu \phi_{s-} + m_s (\bar{\phi}_{s+} \phi_{s+} - \bar{\phi}_{s-} \phi_{s-}) \}, \end{aligned} \quad (2)$$

where $D_\mu \equiv \partial_\mu + ieA_\mu$, $\bar{\psi} \equiv \psi^\dagger \sigma_3$, and m_r and m_s are masses of regulator fields. Of course the mass term of original two-component spinor ψ breaks the parity invariance. Our notations used in this note are the followings;

$$\begin{aligned}\gamma_\mu\gamma_\nu + \gamma_\nu\gamma_\mu &= 2g_{\mu\nu}, \text{diag}[g_{\mu\nu}] = (+1, -1, -1), \\ \gamma_0 &= \sigma_3, \gamma_1 = i\sigma_1, \gamma_2 = i\sigma_2,\end{aligned}\tag{3}$$

$$\begin{aligned}\text{Tr}[\gamma_\mu\gamma_\nu\gamma_\rho\gamma_\sigma] &= 2(g_{\mu\nu}g_{\rho\sigma} + g_{\mu\sigma}g_{\nu\rho} - g_{\mu\rho}g_{\nu\sigma}), \\ \text{Tr}[\gamma_\mu\gamma_\nu\gamma_\rho] &= 2i\varepsilon_{\mu\nu\rho}, \varepsilon_{012} = -1,\end{aligned}\tag{4}$$

where σ 's are Pauli matrices.

Now let us calculate the one loop quantum correction to the regularized photon vacuum polarization tensor $\Pi_{\mu\nu}^M(p)$.

$$\begin{aligned}\Pi_{\mu\nu}^M(p) &= -e^2 \int \frac{d^3k}{(2\pi)^3} \{ \text{Tr}[\gamma_\mu \frac{\gamma^\rho k_\rho - m}{k^2 - m^2} \gamma_\nu \frac{\gamma^\sigma (k+p)_\sigma - m}{(k+p)^2 - m^2}] \\ &\quad + \sum_r \text{Tr}[\gamma_\mu \frac{\gamma^\rho k_\rho - m_r}{k^2 - m_r^2} \gamma_\nu \frac{\gamma^\sigma (k+p)_\sigma - m_r}{(k+p)^2 - m_r^2} \\ &\quad + \gamma_\mu \frac{\gamma^\rho k_\rho + m_r}{k^2 - m_r^2} \gamma_\nu \frac{\gamma^\sigma (k+p)_\sigma + m_r}{(k+p)^2 - m_r^2}] \\ &\quad - \sum_s \text{Tr}[\gamma_\mu \frac{\gamma^\rho k_\rho - m_s}{k^2 - m_s^2} \gamma_\nu \frac{\gamma^\sigma (k+p)_\sigma - m_s}{(k+p)^2 - m_s^2} \\ &\quad + \gamma_\mu \frac{\gamma^\rho k_\rho + m_s}{k^2 - m_s^2} \gamma_\nu \frac{\gamma^\sigma (k+p)_\sigma + m_s}{(k+p)^2 - m_s^2}] \}.\end{aligned}\tag{5}$$

We can see that the contributions of Pauli-Villars regulator fields to the parity odd part (one which contains terms with three γ 's) of $\Pi_{\mu\nu}^M(p)$ automatically cancel and only the contribution of the original spinor field remains. With the aid of the trace formulae of 3-D γ -matrices and Feynmann parameter formulae, after some manipulations and rearrangements, Eq.(5) yields

$$\begin{aligned}\Pi_{\mu\nu}^M(p) &= -2e^2 \int_0^1 dx \int \frac{d^3k}{(2\pi)^3} \{ [\frac{2k_\mu k_\nu}{(k^2 + p^2 x(1-x) - m^2)^2} - \frac{2k_\mu k_\nu}{(k^2 + p^2 x(1-x))^2}] \\ &\quad - [\frac{g_{\mu\nu}}{k^2 + p^2 x(1-x) - m^2} - \frac{g_{\mu\nu}}{k^2 + p^2 x(1-x)}] \\ &\quad + 2k_\mu k_\nu \sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{(k^2 + p^2 x(1-x) - m_n^2)^2} \end{aligned}$$

$$\begin{aligned}
& -g_{\mu\nu} \sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{k^2 + p^2 x(1-x) - m_n^2} \\
& -4x(1-x)(p_\mu p_\nu - p^2 g_{\mu\nu}) \sum_{n=1}^{+\infty} \frac{(-1)^n}{(k^2 + p^2 x(1-x) - m_n^2)^2} \\
& -2x(1-x)(p_\mu p_\nu - p^2 g_{\mu\nu}) \frac{1}{(k^2 + p^2 x(1-x) - m^2)^2} \} \\
& + 2ime^2 \varepsilon_{\mu\nu\rho} p^\rho \int_0^1 dx \int \frac{d^3 k}{(2\pi)^3} \frac{1}{(k^2 + p^2 x(1-x) - m^2)^2}, \tag{6}
\end{aligned}$$

where even $n(= 2r)$ and odd $n(= 2s - 1)$ correspond to contributions from fermionic and bosonic regulator fields, respectively. We have also made the replacement of momentum integration variable; $k_\mu \rightarrow k_\mu - p_\mu x(1-x)$.

The integrals of first and second lines in Eq.(6) are convergent individually and the explicit calculation shows that they cancel each other as they must do. We note that $\int d^3 k k_\mu k_\nu f(k^2) = -\frac{i}{3} g_{\mu\nu} \int d^3 \bar{k} \bar{k}^2 f(-\bar{k}^2)$, where $\bar{k}_\mu$ is the Euclidian three momentum and $f(k^2)$ is any function with suitable analytic properties on the complex energy plane. For notational convenience we define the function $F(\bar{k}^2 + \bar{p}^2 x(1-x)) \equiv \sum_{n=-\infty}^{\infty} \frac{(-1)^n}{\bar{k}^2 + \bar{p}^2 x(1-x) + m_n^2}$, where $\bar{p}^2 \equiv -p^2$. In the following let us assume the mass of regulator fields is given by $m_n = Mn$, although it may not be essential for our discussion. Then the infinite series in Eq.(6) can be explicitly calculated, with the aid of the following formulae;

$$f(a^2) = \sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{a^2 + n^2} = \frac{\pi}{a} \frac{1}{\sinh a\pi}, \tag{7}$$

$$f'(a^2) = \frac{\partial}{\partial a^2} f(a^2) = - \sum_{n=-\infty}^{+\infty} \frac{(-1)^n}{[a^2 + n^2]^2}, \tag{8}$$

$$\sum_{n=1}^{+\infty} \frac{(-1)^n}{a^2 + n^2} = -\frac{1}{2a^2} + \frac{\pi}{2a} \frac{1}{\sinh a\pi}. \tag{9}$$

Clearly $f(a^2)$ exponentially decreases at large a and this ensures integrals involving $f(a^2)$ finite.

The result of the calculation is

$$\Pi_{\mu\nu}^M(p) = -i \frac{e^2}{\pi^2} g_{\mu\nu} \int_0^1 dx \int_0^\infty d\bar{k} \left[\frac{1}{3} \bar{k}^3 \frac{dF(\bar{k}^2 + \bar{p}^2 x(1-x))}{d\bar{k}} + \bar{k}^2 F(\bar{k}^2 + \bar{p}^2 x(1-x)) \right]$$

$$\begin{aligned}
& +i\frac{e^2}{\pi^2 M}(p_\mu p_\nu - p^2 g_{\mu\nu}) \int_0^1 dx x(1-x) \int_0^\infty d\bar{k} \left\{ -\frac{1}{\bar{k}^2 + \frac{\bar{p}^2 x(1-x)}{M^2}} \right. \\
& \left. + \frac{\pi}{\bar{k}^2 + \frac{\bar{p}^2 x(1-x)}{M^2}} \frac{1}{\sinh \pi \sqrt{\bar{k}^2 + \frac{\bar{p}^2 x(1-x)}{M^2}}} \right\} \\
& +i(p_\mu p_\nu - p^2 g_{\mu\nu}) \frac{e^2}{8\pi \bar{p}^2} \left[2m + \frac{\bar{p}^2 - 4m^2}{\bar{p}} \arcsin \sqrt{\frac{\bar{p}^2}{\bar{p}^2 + 4m^2}} \right] \\
& +i^2 \epsilon_{\mu\nu\rho} p^\rho \frac{e^2 m}{2\pi \bar{p}} \arcsin \sqrt{\frac{\bar{p}^2}{\bar{p}^2 + 4m^2}}, \tag{10}
\end{aligned}$$

The first integral in Eq.(10) is identically zero by partial integration. Therefore the gauge invariance for the regularized photon vacuum polarization is indeed maintained independently on the choice of regulator masses m_n . The second integral vanishes as the regulator mass M goes to infinity, since this integral is convergent both in ultraviolet and infrared regions even when $M \rightarrow \infty$. Finally we obtain the following result for the photon vacuum polarization tensor $\Pi_{\mu\nu}(p) \equiv \lim_{M \rightarrow \infty} \Pi_{\mu\nu}^M(p)$.

$$\begin{aligned}
\Pi_{\mu\nu}(p) &= i(p_\mu p_\nu - p^2 g_{\mu\nu}) \frac{e^2}{8\pi \bar{p}^2} \left[2m + \frac{\bar{p}^2 - 4m^2}{\bar{p}} \arcsin \sqrt{\frac{\bar{p}^2}{\bar{p}^2 + 4m^2}} \right] \\
&+ i\epsilon_{\mu\nu\rho} p^\rho \frac{e^2 m}{2\pi \bar{p}} \arcsin \sqrt{\frac{\bar{p}^2}{\bar{p}^2 + 4m^2}}. \tag{11}
\end{aligned}$$

The asymptotic behaviours of $\Pi_{\mu\nu}(p)$ are, as $p \rightarrow 0$ provided $m \neq 0$

$$\Pi_{\mu\nu}(p) \rightarrow i(p_\mu p_\nu - p^2 g_{\mu\nu}) \frac{e^2}{12\pi m} + i^2 \epsilon_{\mu\nu\rho} p^\rho \frac{e^2}{4\pi}, \tag{12}$$

and as $m \rightarrow 0$ provided $p \neq 0$

$$\Pi_{\mu\nu}(p) \rightarrow i(p_\mu p_\nu - p^2 g_{\mu\nu}) \frac{e^2}{16\pi \bar{p}} + i^2 \epsilon_{\mu\nu\rho} p^\rho \frac{e^2 m}{4\bar{p}}. \tag{13}$$

Now it is shown that a parity invariant Pauli-Villars regularization method indeed exists and gives the same results as those obtained by the analytic and dimensional regularization methods. Therefore at least for the perturbative induction of Chern-Simons term in the 3-D quantum electrodynamics three regularization methods lead to the same result.

Finally let us discuss briefly on some remaining problems. Firstly higher-loop correction should be calculated. Secondly the extension to the non-Abelian gauge theories where the new divergent integrals from gauge field loops appear and cause new problems. In Refs.[5] and [7] it was discussed that the gauge invariant regularizations induce the Chern-Simons term (i.e. shift the coefficient of the Chern-Simons term). However the regularization methods in Refs.[5] and [7] break the parity invariance explicitly. Therefore it is an interesting problem to investigate whether any regularization method invariant under both gauge and parity transformations exists for the non-Abelian gauge theories. It seems sufficient to introduce a infinite number of parity doublets of massive vector fields as Pauli-Villars regulators. If this is indeed the case it is expected that no Chern-Simons term is induced even in non-Abelian gauge theories. These problems will be discussed in a subsequent paper.

The present authour would like to thank Toru Ebihara for informing him the work of Frolov and Slavnov.

References

- [1] R. Jackiw and S. Templeton, Phys. Rev. D23, (1981) 2291;
J. F. Schonberg, Nucl. Phys. B185, (1981) 157.
- [2] R. B. Laughlin, Science 242, (1988) 525;
A. L. Fettes, C. B. Hanna and R. B. Laughlin, Phys. Rev. B39, (1989) 9679;
Y. H. Chen, B. I. Halperin, F. Wilczek and E. Witten, Int. J. Mod. Phys B3, (1989) 1001;
N. Dorey and N. E. Mavromatos, Nucl. Phys. B386, (1992) 614;
D. Sénéchal, Phys. Letters B297, (1992) 138.
- [3] E. Witten, Commun. Math. Phys. 121, (1989) 351;
A. P. Polychronakos, Ann. Phys. 203, (1990) 231.
- [4] S..Deser, R. Jackiw and S. Templeton, Ann. Phys. (NY) 140, (1982) 372;
A. N. Redlich, Phys. Rev. D29, (1984) 2366;
Y. -C. Kao and M. Suzuki, Phys. Rev. D31, (1985) 2137.
- [5] L. Alvarez-Gaumé, J. M. F. Labastida and A. V. Ramallo, Nucl. Phys. B334, (1990) 103.
- [6] B. M. Pimentel, A. T. Suzuki and J. L. Tomazelli, Intern. J. Mod. Phys. A7, (1992) 5307.
- [7] C. P. Martin, Phys. Letters B241, (1990) 513.
- [8] R. Delbourgo and A. B. Waites, Phys. Letters B300, (1993) 241.
- [9] S. A. Frolov and A. A. Slavnov, Phys. Letters B309, (1993) 344; Nucl. Phys. B411, (1994) 647.
- [10] R. Narayanan and H. Neugeberger, Phys. Letters B302, (1993) 62;
S. Aoki and Y. Kikukawa, Mod. Phys. Letters A8, (1993) 3517.